

Cathelicidin LL-37 Promotes or Inhibits Cancer Cell Stemness Depending on the Tumor Origin

Guilherme Tude Coelho Neto¹, Thais Martins de Lima¹, Hermes Vieira Barbeiro¹, Roger Chammas², Marcel Cerqueira César Machado¹ and Fabiano Pinheiro da Silva¹✉

1. Emergency Medicine Department, University of Sao Paulo, Sao Paulo, Brazil
2. Oncology Department, University of Sao Paulo, Sao Paulo, Brazil

✉ Corresponding author: Fabiano Pinheiro da Silva, M.D., Ph.D. Faculdade de Medicina da Universidade de São Paulo, Laboratório de Emergências Clínicas (LIM-51), Av. Dr. Arnaldo, 455 sala 3189, CEP 01246-000, São Paulo - SP, Brazil. Phone: +55 11 3061 8480; Fax: +55 11 3061 8480; E-mail: pinheirofabiano@hotmail.com

© Ivyspring International Publisher. Reproduction is permitted for personal, noncommercial use, provided that the article is in whole, unmodified, and properly cited. See <http://ivyspring.com/terms> for terms and conditions.

Abstract

Antimicrobial peptides play critical protective roles in a range of human diseases, including cancer. Multiple studies have demonstrated functions—such as proliferation, angiogenesis, apoptosis and immunomodulation—of these peptides in crucial cancer pathways. We investigated the role of the antimicrobial peptide LL-37 on stemness in breast cancer (SKBR3) and melanoma cells (A375). PCR array analysis of differential gene expression in SKBR3 and A375 cancer cell lines downregulated for LL-37 expression by siRNA revealed downregulation of genes related to stemness, including telomerase reverse transcriptase, forkhead box D3 and undifferentiated embryonic cell transcription factor 1, remarkably in breast cancer cells. Furthermore, SKBR3 cells knocked down for LL-37 expression showed a decreased production of oncospheres in comparison with negative controls, while A375 cells exhibited increased production. Taken collectively, our findings indicate a role for LL-37 in cancer cell stemness depending on the cell type.

Key words: LL-37, cancer, stemness, pluripotency, self-renewal

Introduction

Antimicrobial peptides play crucial roles in critical molecular pathways in cancer, such as cell proliferation, epithelial cell migration, angiogenesis promotion, induction of apoptosis and immunomodulation (1).

The effects of the antimicrobial peptide LL-37 in cancer remain unclear. While LL-37 acts as a positive regulator of ovarian, breast, melanoma and lung cancer progression, it also suppresses colorectal and gastric cancer cell growth (2), indicating that its effects are tumor-specific.

Recently, the concept of clonal tumor evolution has been challenged by the observation of cancer stem

cells (CSCs) in a variety of tumors. CSCs possess increased invasive and metastatic capabilities and render tumors more resistant to several microenvironmental stresses, including the action of several anti-cancer drugs (3).

Here we investigated the effects of LL-37 on stemness in both breast cancer and melanoma cells. We performed array analysis to examine the expression of 84 genes related to DNA damage in wild-type and LL-37-knockdown cancer cells.

Material and Methods

The study protocol was approved by the

Hospital das Clinicas Ethical Committee, protocol 034/14.

Cell culture.

Immortalized human breast cancer cells (SKBR-3) and skin malignant melanoma cells (A375) were used in this study. Cells were maintained according to the guidelines of the ATCC (American Type Culture Collection).

Real-time PCR.

RNA was extracted from cultured cells using TRIZOL[®] protocol. RNA was quantified using NanoVeu (GE Healthcare) systems and RT-PCR was performed using the StepOne SuperScript[®] III (Applied Biosystems) protocol as provided by the manufacturer. Beta-2 microglobulin (B2M) gene was used as an internal control; primers were as follows: GAT GAG TAT GCC TGC CGT TGC, and CAA TCC AAA TGC GGC ATC T. The reactions were performed in a StepOne[™] system (Applied Biosystems) at 50°C for 10 min, 95°C for 5 min and then 95°C for 15 s followed by 60°C for 30 s, and 72°C for 30 s for 40 cycles. Quantification was performed by 2^{-ΔΔCT} method.

LL-37 gene silencing.

Cells were plated at 2.5×10⁵ cells per well in a 6-well plate overnight. LL-37 Silencer Selected Pre-designed short interfering RNA (siRNA) or negative scramble siRNA (Ambion[®]) (10 nM each) was combined with 5 μL of Lipofectamine[™] RNAiMAX reagent for 20 min. Opti-MEM[®] I Reduced Serum Medium (Invitrogen) was added to a final volume of 2.5 mL per well after cells were rinsed with PBS. After 24 h (SKBR3) or 48 h (A375), experimental assays were performed.

PCR array.

Total RNA was converted into cDNA using the RT² First Strand Kit (SABiosciences, Frederick, USA) and cDNA was then combined with the RT² SYBR Green qPCR Master Mix (SABiosciences). Each sample was added to 24 Human DNA Damage PCR arrays (Qiagen, USA) according to the StepOne equipment protocol (Applied Biosystems). PCR-array data analysis was performed at the manufacturer's website (<http://www.sabiosciences.com/pcrarraydataanalysis.php>).

Sphere-forming assays.

The assay was performed as previously reported (4, 5). Briefly, after siRNA treatment, cells were detached and a single cell suspension was obtained

after passing cells through 25 G needles. Cells (1.0×10⁵) were plated in their respective cell culture media containing B27 supplement and rEGF (100 ng/ml; Sigma Aldrich, Poole, UK; E-9644). After 5 days, the number of spheres that were greater than 50 μm in diameter were counted and sphere forming efficiency (%) was determined.

Statistical analysis.

Results were analyzed using Mann-Whitney test and are shown as mean ± standard deviation. A p-value < 0.05 was considered significant.

Results

LL-37 production upregulates several pathways related to stemness.

We next analyzed differential gene expression in cancer cell lines downregulated for LL-37 expression by siRNA. Knockdown efficiency of LL-37 expression was more than 90%, as evaluated by qPCR (data not shown). PCR arrays showed a downregulation of several genes related to stemness, especially in SKBR3, for LL-37 compared with control cells (Tables 1 and 2).

LL-37 knockdown cells show decreased production of oncospheres in breast cancer cells and increased production in melanoma cells.

We next analyzed the production of cancer-derived extracellular vesicles, oncospheres, as a hallmark of stemness in cancer cell lines depleted for LL-37 expression. Our results showed that SKBR3 cells knocked down for LL-37 expression produced a decreased number of oncospheres compared with negative controls (Figure 1A), while A375 produced an increased number of oncospheres compared with negative controls (Figure 1B).




Figure 1. Oncosphere-forming efficiency in SKBR3 (A) and A375 (B) cell lines depleted for LL-37 expression by siRNA.

Table 1. Genes upregulated in SKBR3 breast cancer cells compared with SKBR3 cells transfected with LL-37 siRNA treatment. Genes related to stemness are in bold.

Symbol	Gene Name	Fold Change	p-value
KAT2A	K(lysine) acetyltransferase 2A	1.6337	0.008484
COL2A1	Collagen, type II, alpha 1	1.8903	0.012853
GDF3	Growth differentiation factor 3	1.8402	0.014975
HNF4A	Hepatocyte nuclear factor 4, alpha	1.9064	0.016682
TERT	Telomerase reverse transcriptase	2.2468	0.018056
HAND1	Heart and neural crest derivatives expressed 1	2.1076	0.02039
HSPA9	Heat shock 70kDa protein 9 (mortalin)	3.7997	0.020459
SOX15	SRY (sex determining region Y)-box 15	2.1014	0.02309
NAT1	N-acetyltransferase 1 (arylamine N-acetyltransferase)	26.925	0.026202
ALPL	Alkaline phosphatase, liver/bone/kidney	1.838	0.026346
PARD6A	Par-6 partitioning defective 6 homolog alpha (C. elegans)	14.2326	0.026805
OLIG2	Oligodendrocyte lineage transcription factor 2	1.7598	0.031648
FOXD3	Forkhead box D3	2.6502	0.03305
MYBL2	V-myb myeloblastosis viral oncogene homolog (avian)-like 2	1.6863	0.035958
LIN28A	Lin-28 homolog A (C. elegans)	1.8108	0.036787
TP53	Tumor protein p53	1.9757	0.040021
FGF2	Fibroblast growth factor 2 (basic)	2.0355	0.044985
NES	Nestin	1.8498	0.047692
UTF1	Undifferentiated embryonic cell transcription factor 1	5.2139	0.048209

Discussion

Self-renewal and pluripotency are the key characteristics of stem cells. Here, our findings suggest that LL-37 regulates stemness. Cancer stem cells (CSCs) are a considerable clinical problem, since they are highly resistant to radiation and chemotherapy (6). The mechanism of resistance is not fully understood, but enhanced DNA repair capacities and low intracellular reactive oxidative species concentrations are implicated (7). CSCs also proliferate more slowly than non-stem carcinoma cells, circulate in the bloodstream (8) and can lead to tumor metastasis and relapse. Definitive markers of CSCs do not exist, but our results demonstrated many interesting results in SKBR3 and A375 wild-type cells, when compared with LL-37 knockdown cells. Nestin, for example, is an important marker of CSCs, regulates proliferation, migration and invasion of cancer cells (9) correlating to a worse prognosis (10).

Table 2. Genes upregulated in A375 melanoma cells compared with A375 cells transfected with LL-37 siRNA treatment. Genes related to stemness are in bold.

Symbol	Gene Name	Fold Change	p-value
LIN28A	Lin-28 homolog A (C. elegans)	1.5252	0
TCF3	Transcription factor 3 (E2A immunoglobulin enhancer binding factors E12/E47)	2.9868	0
RUNX2	Runt-related transcription factor 2	1.7998	0.000007
CCNA2	Cyclin A2	1.3786	0.000011
FOXA2	Forkhead box A2	1.5997	0.000018
CD34	CD34 molecule	1.3355	0.000025
BGLAP	Bone gamma-carboxyglutamate (gla) protein	1.5147	0.000044
CDH2	Cadherin 2, type 1, N-cadherin (neuronal)	1.6844	0.000044
FABP7	Fatty acid binding protein 7, brain	1.6157	0.000056
CCNE1	Cyclin E1	1.472	0.000063
OLIG2	Oligodendrocyte lineage transcription factor 2	1.5713	0.000086
NCAM1	Neural cell adhesion molecule 1	1.2164	0.00013
REST	RE1-silencing transcription factor	1.3996	0.000231
TUBB3	Tubulin, beta 3	1.4953	0.000248
CDC42	Cell division cycle 42 (GTP binding protein, 25kDa)	1.5889	0.00025
KRT15	Keratin 15	1.469	0.000278
GATA2	GATA binding protein 2	1.5547	0.000372
ALPL	Alkaline phosphatase, liver/bone/kidney	1.4816	0.000379
GDF3	Growth differentiation factor 3	1.7102	0.000422
TBX3	T-box 3	1.5775	0.000669
AICDA	Activation-induced cytidine deaminase	1.3863	0.001057
RUNX1	Runt-related transcription factor 1	1.2238	0.001949
KLF4	Kruppel-like factor 4 (gut)	1.1139	0.00216
FGF2	Fibroblast growth factor 2 (basic)	1.4219	0.002267
NAT1	N-acetyltransferase 1 (arylamine N-acetyltransferase)	1.3888	0.002528
LEFTY2	Left-right determination factor 2	1.3489	0.003161
MYCN	V-myc myelocytomatosis viral related oncogene, neuroblastoma derived (avian)	1.1672	0.003271
HAND1	Heart and neural crest derivatives expressed 1	1.2117	0.003824
ZFP42	Zinc finger protein 42 homolog (mouse)	1.2421	0.004222
HSPA9	Heat shock 70kDa protein 9 (mortalin)	1.4029	0.005582
FGFR1	Fibroblast growth factor receptor 1	1.3626	0.005687
FGF4	Fibroblast growth factor 4	1.1789	0.006383
NODAL	Nodal homolog (mouse)	1.6944	0.0064
ALDH1A1	Aldehyde dehydrogenase 1 family, member A1	1.2544	0.007374
KAT2A	K(lysine) acetyltransferase 2A	1.2006	0.007585
PAX6	Paired box 6	1.1578	0.008689
DNMT3B	DNA (cytosine-5-)-methyltransferase 3 beta	1.2005	0.011056
HNF4A	Hepatocyte nuclear factor 4, alpha	1.4146	0.011275
CDK1	Cyclin-dependent kinase 1	1.1471	0.012381
COL1A1	Collagen, type I, alpha 1	1.1761	0.015877
LEFTY1	Left-right determination factor 1	1.226	0.015976
DPPA3	Developmental pluripotency associated 2	1.3138	0.024156
NR5A2	Nuclear receptor subfamily 5, group A, member 2	1.1412	0.026799
EP300	E1A binding protein p300	1.3847	0.029084
KAT7	K(lysine) acetyltransferase 7	1.967	0.029885
PARD6A	Par-6 partitioning defective 6 homolog alpha (C. elegans)	1.2703	0.033506
POU5F1	POU class 5 homeobox 1	1.1131	0.042862
ESRRB	Estrogen-related receptor beta	1.2021	0.044154

Telomerase reverse transcriptase, forkhead box D3 (FOXD3) and undifferentiated embryonic cell transcription factor 1 (UTF1) are other genes related to

stemness, so we hypothesized that LL-37 should be important to maintain stem cell identity in breast cancer cells (**Table 1**). Upregulation of telomerase is a prerequisite for cellular immortalization and has been associated with stemness in various human cancers (11). FOXD3 induces cancer progression by epithelial-mesenchymal transition (12) and UTF1 increases stem cells reprogramming to pluripotency (13).

The results were not so evident in A375 melanoma cells, but the presence of Runt-related transcription factors 1 and 2 (14-16), Cadherin 2 (17, 18), Nodal homolog (mouse) (19) and Aldehyde dehydrogenase 1 (family member A1) pointed to the same direction (20) (**Table 2**). However, the production of oncospheres in A375 cells put in evidence that LL-37 has opposing effects on cancer cell stemness depending on the cell type.

Conclusion

Emerging evidence highlights the role of antimicrobial peptides in non-infectious diseases, such as cancer. The mechanisms triggered by antimicrobial peptides are broad and lead to unexpected cell responses. Here, we show that the antimicrobial peptide LL-37 is implicated in cancer stemness. Further research needs to be directed to better clarify this phenomenon and the role of other antimicrobial peptides in this scenario.

Acknowledgement

This work was supported by FAPESP, the Sao Paulo Research Foundation (grant # 2015/00892-4).

Competing Interests

The authors have declared that no competing interest exists.

References

1. Piktel E, Niemirowicz K, Wnorowska U, Watek M, Wollny T, Gluszek K, et al. The Role of Cathelicidin LL-37 in Cancer Development. *Archivum immunologiae et therapiae experimentalis*. 2016;64(1):33-46.
2. Pinheiro da Silva F, Machado MC. Antimicrobial peptides: clinical relevance and therapeutic implications. *Peptides*. 2012;36(2):308-14.
3. Cojoc M, Mabert K, Muders MH, Dubrovska A. A role for cancer stem cells in therapy resistance: cellular and molecular mechanisms. *Seminars in cancer biology*. 2015;31:16-27.
4. Shaw FL, Harrison H, Spence K, Ablett MP, Simoes BM, Farnie G, et al. A detailed mammosphere assay protocol for the quantification of breast stem cell activity. *Journal of mammary gland biology and neoplasia*. 2012;17(2):111-7.
5. Sadovska L, Eglitis J, Line A. Extracellular Vesicles as Biomarkers and Therapeutic Targets in Breast Cancer. *Anticancer research*. 2015;35(12):6379-90.
6. Butof R, Dubrovska A, Baumann M. Clinical perspectives of cancer stem cell research in radiation oncology. *Radiotherapy and oncology : journal of the European Society for Therapeutic Radiology and Oncology*. 2013;108(3):388-96.

7. Skvortsova I, Debbage P, Kumar V, Skvortsov S. Radiation resistance: Cancer stem cells (CSCs) and their enigmatic pro-survival signaling. *Seminars in cancer biology*. 2015;35:39-44.
8. Yang MH, Imrali A, Heesch C. Circulating cancer stem cells: the importance to select. *Chinese journal of cancer research = Chung-kuo yen cheng yen chiu*. 2015;27(5):437-49.
9. Narita K, Matsuda Y, Seike M, Naito Z, Gemma A, Ishiwata T. Nestin regulates proliferation, migration, invasion and stemness of lung adenocarcinoma. *International journal of oncology*. 2014;44(4):1118-30.
10. Ishiwata T, Matsuda Y, Naito Z. Nestin in gastrointestinal and other cancers: effects on cells and tumor angiogenesis. *World journal of gastroenterology*. 2011;17(4):409-18.
11. Terali K, Yilmazer A. New surprises from an old favourite: The emergence of telomerase as a key player in the regulation of cancer stemness. *Biochimie*. 2016;121:170-8.
12. Chu TL, Zhao HM, Li Y, Chen AX, Sun X, Ge J. FoxD3 deficiency promotes breast cancer progression by induction of epithelial-mesenchymal transition. *Biochemical and biophysical research communications*. 2014;446(2):580-4.
13. Galonska C, Smith ZD, Meissner A. In Vivo and in vitro dynamics of undifferentiated embryonic cell transcription factor 1. *Stem cell reports*. 2014;2(3):245-52.
14. Samokhvalov IM. A long way to stemness. *Cell cycle*. 2012;11(16):2965-6.
15. Niu DF, Kondo T, Nakazawa T, Oishi N, Kawasaki T, Mochizuki K, et al. Transcription factor Runx2 is a regulator of epithelial-mesenchymal transition and invasion in thyroid carcinomas. *Laboratory investigation; a journal of technical methods and pathology*. 2012;92(8):1181-90.
16. Vasuri F, Resta L, Fittipaldi S, Malvi D, Pasquinelli G. RUNX-1 and CD44 as markers of resident stem cell derivation in undifferentiated intimal sarcoma of pulmonary artery. *Histopathology*. 2012;61(4):737-43.
17. Pieters T, van Roy F. Role of cell-cell adhesion complexes in embryonic stem cell biology. *Journal of cell science*. 2014;127(Pt 12):2603-13.
18. Farahani E, Patra HK, Jangamreddy JR, Rashedi J, Kawalec M, Rao Pariti RK, et al. Cell adhesion molecules and their relation to (cancer) cell stemness. *Carcinogenesis*. 2014;35(4):747-59.
19. Stefanidis K, Pergialiotis V, Christakis D, Loutradis D, Antsaklis A. Nodal, Nanog, DAZL and SMAD gene expression in human amniotic fluid stem cells. *Journal of stem cells*. 2013;8(1):17-23.
20. Khorrami S, Zavarani Hosseini A, Mowla SJ, Malekzadeh R. Verification of ALDH Activity as a Biomarker in Colon Cancer Stem Cells-Derived HT-29 Cell Line. *Iranian journal of cancer prevention*. 2015;8(5):e3446.